

Neurologin1/2/3/4

Cat.No. 129 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: yes ICC: not recommended IHC: not recommended IHC-P: not tested yet
Clone	4F9
Subtype	IgG2a (κ light chain)
Immunogen	Recombinant protein corresponding to AA 1 to 695 from rat Neurologin1 (UniProt Id: Q62765)
Reactivity	Reacts with: rat (Q62765, Q62888, Q62889,), mouse (Q99K10, Q69ZK9, Q8BYM5, B0F2B4). Other species not tested yet.
Specificity	Recognizes rat and mouse neurologins 1-3 and mouse neurologin 4. K.O. validated
Remarks	ELISA: The ELISA-protocol for membrane proteins is required.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Neurologins form a family of postsynaptic cell surface molecules that interact with β-neurexins. They are 110-120 kDa polypeptides with homology to acetylcholine esterase. Neurologin1 and neurologin3 are specifically localized to post-synaptic densities of excitatory synapses whereas neurologin2 is found exclusively on inhibitory synapses. Mutations in neurologin3 and neurologin4 have been implicated with a rare, heritable form of autism.

Selected References for 129 011

The synaptic proteins neurexins and neurologins are widely expressed in the vascular system and contribute to its functions. Bottos A, Destro E, Rissone A, Graziano S, Cordara G, Assenzio B, Cera MR, Mascia L, Bussolino F, Arese M Proceedings of the National Academy of Sciences of the United States of America (2009) 10649: 20782-7. . **IHC, WB, IP; tested species: chicken**

Synapsin-dependent reserve pool of synaptic vesicles supports replenishment of the readily releasable pool under intense synaptic transmission.

Vasileva M, Horstmann H, Geumann C, Gitler D, Kuner T
The European journal of neuroscience (2012) 368: 3005-20. . **ELISA**

Truncated tau deregulates synaptic markers in rat model for human tauopathy.

Jadhav S, Katina S, Kovac A, Kazmerova Z, Novak M, Zilka N
Frontiers in cellular neuroscience (2015) 9: 24. . **WB**

The association of GPR85 with PSD-95-neurologin complex and autism spectrum disorder: a molecular analysis.

Fujita-Jimbo E, Tanabe Y, Yu Z, Kojima K, Mori M, Li H, Iwamoto S, Yamagata T, Momoi MY, Momoi T
Molecular autism (2015) 6: 17. . **WB**

Neurologin 1 is dynamically exchanged at postsynaptic sites.

Schapitz IU, Behrend B, Pechmann Y, Lappe-Siefke C, Kneussel SJ, Wallace KE, Stempel AV, Buck F, Grant SG, Schweizer M, Schmitz D, et al.

The Journal of neuroscience : the official journal of the Society for Neuroscience (2010) 3038: 12733-44. . **WB; tested species: rat**

Postsynaptic neurologin enhances presynaptic inputs at neuronal nicotinic synapses.

Conroy WG, Nai Q, Ross B, Naughton G, Berg DK
Developmental biology (2007) 3071: 79-91. . **WB**

A mutation linked with autism reveals a common mechanism of endoplasmic reticulum retention for the alpha,beta-hydrolase fold protein family.

De Jaco A, Comoletti D, Kovarik Z, Gaietta G, Radic Z, Lockridge O, Ellisman MH, Taylor P
The Journal of biological chemistry (2006) 28114: 9667-76. . **WB**

Control of excitatory and inhibitory synapse formation by neurologins.

Chih B, Engelman H, Scheiffele P
Science (New York, N.Y.) (2005) 3075713: 1324-8. . **WB**

Selected General References

Neurologin 1 is a postsynaptic cell-adhesion molecule of excitatory synapses.

Song JY et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PubMed:9927700

Activity-dependent validation of excitatory versus inhibitory synapses by neurologin-1 versus neurologin-2.

Chubykin AA et al. Neuron (2007) PubMed:17582332

Dissection of synapse induction by neurologins: effect of a neurologin mutation associated with autism.

Chubykin AA et al. J. Biol. Chem. (2005) PubMed:15797875

Access the online factsheet including applicable protocols at <https://sysy.com/product/129011> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.