

## Gephyrin

**Cat.No. 147 318; Recombinant Guinea pig antibody, 200 µl recombinant IgG supernatant (lyophilized)**

### Data Sheet

|                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reconstitution/<br>Storage | 200 µl purified recombinant IgG supernatant, lyophilized. For <b>reconstitution</b> add 200 µl H <sub>2</sub> O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Applications               | <b>WB:</b> not recommended (see remarks)<br><b>IP:</b> not recommended (see remarks)<br><b>ICC:</b> 1 : 200<br><b>IHC:</b> 1 : 200 (see remarks)<br><b>IHC-P:</b> not recommended<br><b>IHC-Fr:</b> 1 : 200 up to 1 : 500 (see remarks)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Clone                      | GpmAb7a                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Subtype                    | IgG2 (κ light chain)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Immunogen                  | Nativ Protein corresponding to AA 1 to 768 from rat Gephyrin (UniProt Id: Q03555)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Epitop                     | AA 264 to 276 from rat Gephyrin (UniProt Id: Q03555)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Reactivity                 | Reacts with: human (Q9NQX3), rat (Q03555), mouse (Q8BUV3), pig, goldfish, zebrafish.<br>Other species not tested yet.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Specificity                | Specific for the brain specific 93 kDa splice variant phosphorylated at Ser-270. K.O. validated                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Remarks                    | This antibody is a chimeric antibody based on the well known clone mAb7a. The constant regions of the heavy and light chains have been replaced with Guinea pig specific sequences. The antibody can therefore be used with standard anti-Guinea pig secondary reagents. It also carries a Strep-tag at the C-terminus of the heavy chain and has been expressed in mammalian cells.<br><b>WB:</b> Clone 3B11 (cat. no. <a href="#">147 111</a> ) is highly recommended.<br><b>IP:</b> Clone 3B11 (cat. no. <a href="#">147 111</a> ) highly recommended.<br><b>IHC:</b> Antigen retrieval with citrate buffer pH 6 can be applied to improve the signal to noise ratio.<br>Alternatively, the mild fixation protocol according to in <a href="#">Schneider Gasser et al. 2006</a> , can be applied. Clone mAb7a and its recombinant derivatives can cause non-specific nuclear staining, which becomes more pronounced after AGR treatment. The culture supernatants ( <a href="#">147 021</a> , <a href="#">147 318</a> and <a href="#">147 018</a> ) are less prone to this effect and are the preferred choices for IHC experiments.<br><b>IHC-Fr:</b> The following fixatives are possible: 4% formaldehyde/PFA, acetone, methanol-acetone, methanol<br>Signal intensities as follows: PFA = acetone > acetone-methanol > methanol. |

## Background

**Gephyrin** is a bifunctional protein which is essential for both synaptic clustering of inhibitory neurotransmitter receptors in the central nervous system and the biosynthesis of the molybdenum cofactor (MoCo) in peripheral tissues. It co-purifies with the inhibitory glycine receptor (GlyR) and is expressed abundantly in all brain areas which contain synapses.

## Selected References for 147 318

- Necl-4/CADM4 regulates GABAergic synaptic strength on GABAergic inhibitory neurons via ErbB4 activation and prevents neuronal impairments.  
Komaki R, Shiotani H, Kuriu T, Kameyama T, Miyata M, Kedashiro S, Katanazaka K, Nishii S, Chihara N, Matsumoto R, Koebis M, et al.  
Molecular neurobiology (2025) : . . **ICC, IHC; tested species: mouse**
- Semaphorin 3A-mediated perineuronal nets formation incubates depressive-like behaviors in male mice via activating parvalbumin-expressing interneurons.  
Mei YX, Chen HS, Cao QL, Lv WW, Liu Y, Deng SL, Zhou Q, Zhang ZJ, Wang F, Hu ZL, Chen JG, et al.  
Molecular psychiatry (2025) : . . **IHC; tested species: mouse**
- Dual lineage origins contribute to neocortical astrocyte diversity.  
Zhou J, Vitali I, Roig-Puigros S, Javed A, Cantando I, Puglisi M, Bezzi P, Jabaudon D, Mayer C, Bocchi R  
Nature communications (2025) 161: 6992. . **IHC; tested species: mouse**
- Distinct mechanisms control the specific synaptic functions of Neuroligin 1 and Neuroligin 2.  
Wang J, Sudhof T, Wernig M  
EMBO reports (2025) : . . **ICC; tested species: mouse**
- The TMEM132B-GABAA receptor complex controls alcohol actions in the brain.  
Wang G, Peng S, Reyes Mendez M, Karamidas A, Castellano D, Wu K, Han W, Tian Q, Dong L, Li Y, Lu W, et al.  
Cell (2024) 18723: 6649-6668.e35. . **ICC; tested species: mouse**
- Cholecystokinin-expressing interneurons mediated inhibitory transmission and plasticity in basolateral amygdala modulate stress-induced anxiety-like behaviors in mice.  
Fang W, Chen X, He J  
Neurobiology of stress (2024) 33: 100680. . **IHC; tested species: mouse**
- Biphasic Npas4 expression promotes inhibitory plasticity and suppression of fear memory consolidation in mice.  
Brito DVC, Kupke J, Sokolov R, Cambridge S, Both M, Bengtson CP, Rozov A, Oliveira AMM  
Molecular psychiatry (2024) : . . **IHC; tested species: mouse**
- Neuronal activation of Gqα EGL-30/GNAQ late in life rejuvenates cognition across species.  
Stevenson ME, Bieri G, Kaletsky R, St Ange J, Remesal L, Pratt KJB, Zhou S, Weng Y, Murphy CT, Villeda SA  
Cell reports (2023) 429: 113151. . **IHC; tested species: mouse**
- A thalamic-primary auditory cortex circuit mediates resilience to stress.  
Li HY, Zhu MZ, Yuan XR, Guo ZX, Pan YD, Li YQ, Zhu XH  
Cell (2023) 1867: 1352-1368.e18. . **IHC; tested species: mouse**
- Preclinical long-term safety of intraspinal transplantation of human dorsal spinal GABA neural progenitor cells.  
Zheng X, Liu Z, He Z, Xu J, Wang Y, Gong C, Zhang R, Zhang SC, Chen H, Wang W  
iScience (2023) 2611: 108306. . **IHC; tested species: rat**
- Gabrb3 is required for the functional integration of pyramidal neuron subtypes in the somatosensory cortex.  
Babij R, Ferrer C, Donatelle A, Wacks S, Buch AM, Niemeyer JE, Ma H, Duan ZRS, Fetcho RN, Che A, Otsuka T, et al.  
Neuron (2022) : . . **IHC; tested species: mouse**
- Neurotransmitter phenotype switching by spinal excitatory interneurons regulates locomotor recovery after spinal cord injury.  
Bertels H, Vicente-Ortiz G, El Kanbi K, Takeoka A  
Nature neuroscience (2022) 255: 617-629. . **IHC; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/147318> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.