

Veli1/2/3

Cat.No. 184 002; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For reconstitution add 200 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not tested yet ICC: 1 : 1000 IHC: not tested yet IHC-P: not tested yet
Immunogen	Recombinant protein corresponding to AA 1 to 232 from rat Velis1 (UniProt Id: Q9Z250)
Reactivity	Reacts with: human (O14910, Q9HAP6, Q9NUP9), rat (Q9Z250, Q9Z252, Q79210), mouse (Q8JZS0, O88951, O88952). Other species not tested yet.
Specificity	Recognizes all three isoforms veli 1, 2 and 3.
Matching control	184-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

The complex of the multi-adaptor-proteins CASK, Mint and **Veli**, also referred to as **MALS** and **Lin 7**, is conserved across kingdoms and may be involved in localizing synaptic vesicles at synaptic terminals. Three Veli isoforms (1, 2 and 3) with different expression patterns have been described, so far. Veli proteins have also been suggested to participate in regulating the recruitment of neurotransmitter receptors to the post synaptic density (PSD) by interaction with their PDZ domains.

Selected References for 184 002

Functional analysis of CASK transcript variants expressed in human brain.
Tibbe D, Pan YE, Reißner C, Harms FL, Kreienkamp HJ
PloS one (2021) 166: e0253223. . **WB; tested species: human**

Liprin-α proteins are master regulators of human presynapse assembly.
Marcó de la Cruz B, Campos J, Molinaro A, Xie X, Jin G, Wei Z, Acuna C, Sterky FH
Nature neuroscience (2024) : . . **WB; tested species: human**

Regulation of Liprin-α phase separation by CASK is disrupted by a mutation in its CaM kinase domain.
Tibbe D, Ferle P, Krisp C, Nampoothiri S, Mirzaa G, Assaf M, Parikh S, Kutsche K, Kreienkamp HJ
Life science alliance (2022) 510: . . **WB; tested species: human**

Selected General References

Synaptic transmission regulated by a presynaptic MALS/Liprin-alpha protein complex.
Olsen O et al. Curr. Opin. Cell Biol. (2006) PubMed:16504495

Neurotransmitter release regulated by a MALS-liprin-alpha presynaptic complex.
Olsen O et al. J. Cell Biol. (2005) PubMed:16186258

The SNARE motif contributes to rbet1 intracellular targeting and dynamics independently of SNARE interactions.
Joglekar AP et al. J. Biol. Chem. (2003) PubMed:12566453

Access the online factsheet including applicable protocols at <https://sysy.com/product/184002> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.