

m6A

Cat.No. 202 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	Dot blot: 1 : 1000 up to 1 : 10000 IP: yes ICC: not tested yet IHC: not tested yet IHC-P: not tested yet ELISA: suitable for sandwich-ELISA MeRIP: yes
Immunogen	N6-methyladenosine fused to BSA
Reactivity	Reacts with: human, rat, mouse, eukaryotes, prokaryotes. Other species not tested yet.
Specificity	Specific for N6-methyladenosine (m6A) with some cross-reactivity to m6Am.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

m6A (N6-methyladenosine) is a posttranscriptional RNA-modification found throughout all kingdoms, e.g. in vertebrate snRNAs U2, U4, U6, in viral and eukaryotic mRNAs, and in E. coli 16S rRNA. Recent studies have found that mRNA is predominately m6A modified at stop codons and long internal exons, which are conserved between mouse and human. The so-called RNA methylome probably plays an important role in the regulation of gene expression.

In E. coli Dam methylase introduces m6A modifications on the DNA level at the 5'-GATC-3' motif. This allows the cell to differentiate between the parental and the daughter strand during mismatch repair.

Selected References for 202 003

N6-methyladenine DNA Modification in Glioblastoma.

Xie Q, Wu TP, Gimple RC, Li Z, Prager BC, Wu Q, Yu Y, Wang P, Wang Y, Gorkin DU, Zhang C, et al. Cell (2018) : . . . **DOTBLOT, IP, ICC, IHC-P; tested species: human**

Multiplexed profiling facilitates robust m6A quantification at site, gene and sample resolution.

Dierks D, Garcia-Campos MA, Uzonyi A, Safra M, Edelheit S, Rossi A, Sideri T, Varier RA, Brandis A, Stelzer Y, van Werven F, et al. Nature methods (2021) : . . . **IP; tested species: mouse**

Extracellular matrix stiffness reduces DNA 6 ma level to facilitate colorectal cancer progression via disrupting P53 binding to CDKN1A promoter.

Xie SA, Li X, Yin MY, Du F, Zhang ST, Zhu ST

Experimental hematology & oncology (2025) 141: 111. . . **DOTBLOT, IP, IHC, ELISA; tested species: human,mouse**

Colchicine Blocks Abdominal Aortic Aneurysm Development by Maintaining Vascular Smooth Muscle Cell Homeostasis.

Chen M, Yang D, Zhou Y, Yang C, Lin W, Li J, Liu J, Ye J, Huang W, Ma W, Li W, et al.

International journal of biological sciences (2024) 206: 2092-2110. . . **DOTBLOT, IHC, MERIP; tested species: mouse,human**

Epstein-Barr virus suppresses N6-Methyladenosine modification of TLR9 to promote immune evasion.

Zhang X, Li Z, Peng Q, Liu C, Wu Y, Wen Y, Zheng R, Xu C, Tian J, Zheng X, Yan Q, et al.

The Journal of biological chemistry (2024) : 107226. . . **WB, MERIP, DOTBLOT; tested species: human**

Reversible ALKBH5 cytosolic aggregation accelerates cellular senescence.

Chen L, Chen Z, Mo J, Xie Q, Lin W, Zheng H, Zou Y, Chen Y, Bian XW, Zhou Z, Jin G, et al.

Cell death and differentiation (2025) : . . . **DOTBLOT, ICC, MERIP; tested species: human,mouse**

Multichrome encoding-based multiplexed, spatially resolved imaging reveals single-cell RNA epigenetic modifications heterogeneity.

Mao D, Tang X, Zhang R, Hu S, Gou H, Zhang P, Li W, Pan Q, Shen B, Zhu X

Nature communications (2025) 161: 958. . . **ICC, IHC, MERIP; tested species: human,mouse**

METTL3 Promotes Osteogenic Differentiation of Human Periodontal Ligament Stem Cells through IGF2BP1-Mediated Regulation of Runx2 Stability.

Sun X, Meng X, Piao Y, Dong S, Dong Q

International journal of medical sciences (2024) 214: 664-673. . . **DOTBLOT, IP, ICC; tested species: human**

Decoding the interplay between m6A modification and stress granule stability by live-cell imaging.

Li Q, Liu J, Guo L, Zhang Y, Chen Y, Liu H, Cheng H, Deng L, Qiu J, Zhang K, Goh WSS, et al.

Science advances (2024) 1046: eadp5689. . . **DOTBLOT, ICC, MERIP; tested species: human**

METTL3 enhances dentinogenesis differentiation of dental pulp stem cells via increasing GDF6 and STC1 mRNA stability.

Pan Y, Liu Y, Cui D, Yu S, Zhou Y, Zhou X, Du W, Zheng L, Wan M

BMC oral health (2023) 231: 209. . . **IP, DOTBLOT, IHC-P; tested species: mouse**

Downregulation of ALKBH5 rejuvenates aged human mesenchymal stem cells and enhances their therapeutic efficacy in myocardial infarction.

Gao X, Liang X, Liu B, Hong Y, He H, Shen Y, Chen J, Huang X, Hu B, Li W, Li X, et al.

FASEB journal : official publication of the Federation of American Societies for Experimental Biology (2023) 3712: e23294. . . **IP, DOTBLOT, ICC; tested species: human**

Access the online factsheet including applicable protocols at <https://sysy.com/product/202003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.