

Calbindin D28k

Cat.No. 214 318; Recombinant Guinea pig antibody, 50 µg recombinant IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not tested yet ICC: 1 : 500 up to 1 : 2000 IHC: 1 : 500 IHC-P: 1 : 500 up to 1 : 2000
Clone	Gp351C10
Subtype	IgG2 (κ light chain)
Immunogen	Full-length recombinant human Calbindin D28k protein (UniProt Id: P05937)
Reactivity	Reacts with: human (P05937), rat (P07171), mouse (P12658), zebrafish, grasshopper. Other species not tested yet.
Matching control	214-0P
Remarks	This antibody is a chimeric antibody based on the monoclonal mouse antibody 351C10. The constant regions of the heavy and light chains have been replaced with guinea pig specific sequences. Therefore, the antibody can be used with standard anti-guinea pig secondary reagents. The antibody has been expressed in mammalian cells. IHC: Antigen retrieval with citrate buffer pH 6 is tolerated. For optimal results in retina tissue, follow the retina protocol according to Gierke et al. 2023 .

Background

Calbindin D28k (CALB1, D28K or CAB27) and Calretinin (Calbindin D29k) are members of the EF-hand calcium-binding protein family, playing essential roles in intracellular calcium homeostasis, neuroprotection, and signal transduction (1). Calbindin D28k is widely expressed in tissues that experience high calcium influx, such as the brain, sensory organs, endocrine tissues, and the enteric nervous system (ENS). In the central nervous system (CNS), Calbindin D28k is an important marker for specific neuronal populations, particularly in Purkinje cells of the cerebellum and GABAergic and glutamatergic neurons of the cortex (2). Outside the CNS, Calbindin D28k plays crucial roles in non-neuronal tissues. In the retina, it contributes to photoreceptor function and calcium signaling (3). In the pancreas, it is present in α- and β-cells (4). Additionally, it was reported that 31% of myenteric neurons and up to 95% of submucosal neurons in the human ENS express calbindin, suggesting its involvement in both motor and secretory functions (5).

Selected General References

'New' functions for 'old' proteins: the role of the calcium-binding proteins calbindin D-28k, calretinin and parvalbumin, in cerebellar physiology. Studies with knockout mice.
Schwaller B et al. Cerebellum (2002) PubMed:12879963

Calbindin D28k-Immunoreactivity in Human Enteric Neurons.
Zetzmann K et al. Int J Mol Sci (2018) PubMed:29316719

Types and density of calbindin D28k-immunoreactive ganglion cells in mouse retina.
Gu YN et al. Exp Eye Res (2016) PubMed:26874036

Calbindin-D28K immunoreactivity in the human cerebellar cortex.
Flace P et al. Anat Rec (Hoboken) (2014) PubMed:24719368

Expression of the novel epithelial Ca²⁺ channel ECaC1 in rat pancreatic islets.
Janssen SW et al. J Histochem Cytochem (2002) PubMed:12019295

Access the online factsheet including applicable protocols at <https://sysy.com/product/214318> or scan the QR-code.



TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.