

## Syncollin

Cat.No. 225 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

|                            |                                                                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reconstitution/<br>Storage | 100 µg purified IgG, lyophilized. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet. |
| Applications               | <b>WB:</b> 1 : 1000 up to 1 : 5000 (AP staining)<br><b>IP:</b> not tested yet<br><b>ICC:</b> not tested yet<br><b>IHC:</b> not tested yet<br><b>IHC-P:</b> not recommended                                                                                                                                    |
| Clone                      | 87.1                                                                                                                                                                                                                                                                                                          |
| Subtype                    | IgG1 (κ light chain)                                                                                                                                                                                                                                                                                          |
| Immunogen                  | Recombinant protein corresponding to AA 22 to 134 from rat Syncollin (UniProt Id: O35775)                                                                                                                                                                                                                     |
| Reactivity                 | Reacts with: rat (O35775), mouse (Q8VCK7).<br>Other species not tested yet.                                                                                                                                                                                                                                   |

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

The exocrine pancreas is specialized for the synthesis and secretion of digestive enzymes which are stored in zymogen granules. **Syncollin** is predominantly associated with the luminal membrane surface of these organelles. It forms oligomers and can bind to syntaxin 1 and syntaxin 2 in a calcium dependent manner. The exact physiological role has not yet been elucidated.

## Selected References for 225 011

The secretory granule protein syncollin localizes to HL-60 cells and neutrophils.  
Bach JP, Borta H, Ackermann W, Faust F, Borchers O, Schrader M  
The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society (2006) 548: 877-88. . **WB; tested species: rat**

Interaction of syncollin with GP-2, the major membrane protein of pancreatic zymogen granules, and association with lipid microdomains.  
Kalus I, Hodel A, Koch A, Kleene R, Edwardson JM, Schrader M  
The Biochemical journal (2002) 362Pt 2: 433-42. . **WB**

Loss of the zymogen granule protein syncollin affects pancreatic protein synthesis and transport but not secretion.  
Antonin W, Wagner M, Riedel D, Brose N, Jahn R  
Molecular and cellular biology (2002) 225: 1545-54. . **WB; KO verified; tested species: mouse**

Analysis of the association of syncollin with the membrane of the pancreatic zymogen granule.  
An SJ, Hansen NJ, Hodel A, Jahn R, Edwardson JM  
The Journal of biological chemistry (2000) 27515: 11306-11. . **WB**

## Selected General References

The secretory granule protein syncollin localizes to HL-60 cells and neutrophils.  
Bach JP et al. J. Histochem. Cytochem. (2006) PubMed:16517980

Interaction of syncollin with GP-2, the major membrane protein of pancreatic zymogen granules, and association with lipid microdomains.  
Kalus I et al. Biochem. J. (2002) PubMed:11853552

Loss of the zymogen granule protein syncollin affects pancreatic protein synthesis and transport but not secretion.  
Antonin W et al. Mol. Cell. Biol. (2002) PubMed:11839820

Analysis of the association of syncollin with the membrane of the pancreatic zymogen granule.  
An SJ et al. J. Biol. Chem. (2000) PubMed:10753942

Access the online factsheet including applicable protocols  
at <https://sysy.com/product/225011> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.