

Doublecortin

Cat.No. 326 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For reconstitution add 100 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: yes ICC: not recommended IHC: not recommended IHC-P: 1 : 500
Immunogen	Recombinant protein corresponding to AA 271 to 366 from mouse Doublecortin (UniProt Id: Q6PGI2)
Reactivity	Reacts with: rat (Q9ESI7), mouse (Q6PGI2). Other species not tested yet.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Doublecortin or **DCX** seems to be required for initial steps of neuronal dispersion and cortex lamination during cerebral cortex development. Mutations in the gene encoding this protein have been linked to the double cortex syndrome.

Doublecortin is highly expressed in neuronal cells of fetal brain but absent from other tissues. In the adult dentate gyrus it is only expressed in cells contributing to adult neurogenesis.

Selected References for 326 004

Impaired p53-Mediated DNA Damage Response Contributes to Microcephaly in Nijmegen Breakage Syndrome Patient-Derived Cerebral Organoids.

Martins S, Erichsen L, Datsi A, Wruck W, Goering W, Chatzantonaki E, de Amorim VCM, Rossi A, Chrzanowska KH, Adjaye J Cells (2022) 115: . . **IHC; tested species: human**

Stem cell expression of CXCR4 regulates tissue composition in the vomeronasal organ.

Dietz A, Senf K, Neuhaus EM

Journal of cell science (2025) 1381: . . **IHC; tested species: mouse**

Azithromycin preserves adult hippocampal neurogenesis and behavior in a mouse model of sepsis.

Rodríguez-Moreno CB, Cañeque-Rufo H, Flor-García M, Terreros-Roncal J, Moreno-Jiménez EP, Pallas-Bazarra N, Bressa C, Larrosa M, Cofini F, Llorens-Martín M

Brain, behavior, and immunity (2024) 117: 135-148. . **IHC; tested species: mouse**

Prolonged fixation and post-mortem delay impede the study of adult neurogenesis in mice.

Gallardo-Caballero M, Rodríguez-Moreno CB, Álvarez-Méndez L, Terreros-Roncal J, Flor-García M, Moreno-Jiménez EP, Rábano A, Llorens-Martín M

Communications biology (2023) 61: 978. . **IHC; tested species: mouse**

Preclinical long-term safety of intraspinal transplantation of human dorsal spinal GABA neural progenitor cells.

Zheng X, Liu Z, He Z, Xu J, Wang Y, Gong C, Zhang R, Zhang SC, Chen H, Wang W

iScience (2023) 2611: 108306. . **IHC; tested species: rat**

Selected General References

Redirection of doublecortin-positive cell migration by over-expression of the chemokines MCP-1, MIP-1α and GRO-α in the adult rat brain.

Tang SK et al. Neuroscience (2014) PubMed:24361178

Variability of doublecortin-associated dendrite maturation in adult hippocampal neurogenesis is independent of the regulation of precursor cell proliferation.

Plümpe T et al. BMC Neurosci (2006) PubMed:17105671

Distinct roles of doublecortin modulating the microtubule cytoskeleton.

Moores CA et al. EMBO J. (2006) PubMed:16957770

Doublecortin is a developmentally regulated, microtubule-associated protein expressed in migrating and differentiating neurons.

Francis F et al. Neuron (1999) PubMed:10399932

Doublecortin, a brain-specific gene mutated in human X-linked lissencephaly and double cortex syndrome, encodes a putative signaling protein.

Gleeson JG et al. Cell (1998) PubMed:9489700

Access the online factsheet including applicable protocols at <https://sysy.com/product/326004> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.